

# Solar and IP precursors of Geomagnetic storms in SC 23

J. Uwamahoro <sup>(1)</sup>    L. A. McKinnell <sup>(2)</sup>

<sup>(1)</sup>Department of Maths and Physics, University of Rwanda, College of Education



<sup>(2)</sup>South African National Space Agency [SANSA]-Space Science-Hermanus, SA  
**UN/AUSTRIA Symposium on Space Weather Data, Instruments and Models,  
Graz, 16-18/09/2013**



# Outline

- 1 **Introduction**
  - Characteristics of geoeffective solar and IP events
  - Geomagnetic storms
- 2 **Data and methods**
  - Methods of investigation
- 3 **Results and discussion**
  - Statistics of events in SC 23
- 4 **Summary**

# Solar transients: CMEs and Solar Flares




# Coronal Mass Ejections, Geoeffective CMEs

- Transients expulsions of plasma and magnetic field from the Sun
- Produce disturbances in the IP medium leading to phenomena known as **geomagnetic storms**
- GMS are strong perturbations of the Earth atmosphere affecting **space weather** in various ways



# CMEs characteristics

Property	range	average
Speed	$\sim 20 \text{ km/s}$ to $> 3000 \text{ km/s}$	$\sim 470 \text{ km/s}$
Mass	$\sim 10^{12} \text{ g}$ to $> 10^{16} \text{ g}$	$\sim 4 \times 10^{14} \text{ g}$
Kinetic energy	$\sim 10^{27} \text{ erg}$ to $10^{33} \text{ erg}$	$\sim 5 \times 10^{29} \text{ erg}$
Angular width	$< 5^\circ$ to $360^\circ$	$\sim 54^\circ$
Daily occurrence rate	$< 0.5$ to $> 6$ CMEs	Solar min - solar max


# Main sources of geomagnetic storms



# CIRs as sources of recurrent GMS




# Magnetic storms generation





# Space Weather impact



# Properties of Geoeffective CMEs

- 1 **Geoeffectivity**: Their ability to produce geomagnetic storms: In this study,  $Dst \leq -50$  nT.
- 2 **halo CMEs**: appear to surround the occulting disk of the observing coronagraphs
- 3 Generally fast and wide and mostly associated with powerful flares (Class X and M)
- 4 **Full** halo CMEs: apparent width ( $W$ ) of  $360^\circ$
- 5 **partial** halo CMEs: apparent width ( $W$ ) of  $120^\circ \leq W \leq 360^\circ$ .
- 6 But still not very clear what kind of CMEs produce GMS, some halo and front-sided CMEs may not have a geomagnetic impact

# Objectives

- Recent decades, intensive research focusing on estimating the geoeffectiveness of solar phenomena

# Objectives

- A statistical investigation of CMEs and associated solar and IP properties that were probable causes of 229 magnetic storms covering 1996-2006; a full average 11-year solar cycle.

# Objectives

- Comparison of the magnetic storm effectiveness between full and partial halo CMEs


# Selection of geomagnetic storms events

- Selection based on the **Dst** index
- A measure of the H-component of the Earth's MF at low to mid latitude
- A good measure of the ring current
  - 1 Moderate storms (with  $-100 \text{ nT} \leq Dst \leq -50 \text{ nT}$ )
  - 2 Intense storms with  $Dst \leq -100 \text{ nT}$
  - 3 Minor storms ( $Dst > -50 \text{ nT}$ ) **not considered**.
- In total, 244 storm events identified (Jan 1996- Dec 2006).
- Data source on :  
<http://www.ngdc.noaa.gov/stp/geomag/dst.html>.

# IP signatures of geoeffective solar events

- Transport of solar disturbances to near Earth via Solar wind (SW).
- CME structures at 1 AU are known as **ICMEs**.
- ICMEs are geoeffective when associated with negative Z-component of **IMF(Bs)**.
- A table of ICMEs with associated properties and geomagnetic effect by **(Richardson and Cane, 2010)**.  
(see:<http://www.srl.caltech.edu/ACE/ASC/DATA/level3/icme>)
- We produced a similar table based on GMS events (no minor storms involved) which shows additional **92 storm events**
- Data source on **OMNIWEB**:  
<http://www.nssdc.gsfc.nasa/omniweb.html>.

# Geoeffective SW structures associated with the passage of an ICME in IP medium





# Geoeffective properties associated with halo CMEs

- Angular Width (AW) of CMEs as appear in solar coronagraphs: large AW implies higher probability of Earth's impact
- CME speed and association with powerful solar flares
- Surface location of a long duration flare can be considered as the source region of the associated CME
- Considered a range of  $\pm 0.5$  hours to decide the association of halo CME eruption with flare occurrence

# Data source

## Solar data from..

- 1 LASCO/CME data:  
[http://cdaw.gsfc.nasa.gov/CME\\_list](http://cdaw.gsfc.nasa.gov/CME_list)
- 2 GOES data on:  
<http://www.ngdc.noaa.gov/stp/solar/solarflares.html>.

# Methods of investigation

- Identified solar and IP parameters ( $AW_{cmes}$ ,  $V_{cmes}$ ,  $B_S$ ,  $V_{sw}$ ,  $SF_S$ ) suitable enough for analysis of storm efficiency.
- Prediction of storm occurrence at 86% when used in an empirical model, [Uwamahoro et al., 2012](#).

## Method

“A 5-day time window prior to the occurrence of a storm was used to explore probable halo CMEs (and associated solar and IP properties) causes of the subsequent storm. ”

- But one storm event may follow from more than one halo CME; Consider frontside CMEs with other associated geoeffective properties.

# Selected examples from a Table of 244 storm events


Event/ parameters	Event 232	Event 187	Event 195
Date of event	24/08/2005	20/11/2003	04/04/04
Dst(nT)	-216	-422	-112
$B_S$ (nT)	-38.3	-50.9	-7.9
$V_{sw}$ (km/h)	620	553	506
Halo CME	<i>FH</i> : 22/08[01 : 31; 17 : 30]	<i>FH</i> : 18/11[08 : 50]	CIR or SOHO stealth CME..??
$V_{cme}$ (km/h)	1194; 2378	1668	-
Flare/Location	<i>M2.6</i> ; <i>S11 W54</i>	<i>M3.9</i> ; <i>N03E18</i>	-
ICME/Time	24/08 [14:00]	20/11 [10:00]	03/04 [14:00]

# Results and discussion

## Main results

- 1 In total, 244 events [1996-2006]; excluding 15 events of missing SOHO/LASCO data
- 2 July, August, September 1998 and January 1999], hence analysis of **229** storm events
- 3 Identified 84 intense GMS and 145 moderate storms

# Solar and IP precursors of intense vs moderate storms







A= Distribution of sources for moderate storms: (a) CIRs, (b) FH CMEs; (c) PH CMEs.

B= Distribution for intense storms: (a) FH CMEs, (b) PH CMEs (c) CIRs

# Solar and IP precursors of intense vs moderate storms









Storm category	No.of GMS	FH-CME	PH-CME	No CMEs	ICMEs
Intense storms	84	55 [65%]	15 [18%]	14 [17%]	72[86%]
Moderate storms	145	48 [33%]	31 [21%]	66 [46%]	64 [44%]
Total	229	103 [45%]	46 [20%]	80[35%]	136 [59%]

# IP properties for intense vs moderate storms





# Intense vs moderate storms associated properties







# On magnetic storms association with solar flares

<b>Flare class</b>	<b>B-class</b>	<b>C-class</b>	<b>M-class</b>	<b>X-class</b>
Number	9	46	48	23
Percentage	7%	37%	38%	18%



# Multiple halo CME associated storms

- 39 cases identified; mostly lead to intense storms (up to 69%).
- Generally associated with higher average values of  $B_s$ ,  $V_{SW}$  accompanied by ICME at 92%




## Multiple CME storms precursors properties

# Multiple vs one CME driven storm



# Trends in SC 23


- Observed correlation between solar activity (in terms of SSN) and the occurrence of storms.
- Notice a triple peak in both CME and GMS occurrence



SC trends in CMEs and associated GMS.

# Trends in SC 23


- Observed trends by SC phase in terms of GMS, halo CMEs, ICMEs and CIRs.
- Half of all nonhalo CME associated storms accrued in the declining phase



Frequency by SC phase.

# Trends in SC 23

- Frequency occurrence per year in SC 23 of 244 GMS and associated properties.
- Peaks in 1998, 200-2002 and 2005. Exceptional peak in 2003 due to **Halloween storms**.



Frequency per year.

# Main results

## Summary of main results

- 1 During an average 11-year SC, 83% of intense storms were found to be caused by halo CMEs
- 2 CIRs and / or undetected CMEs mostly moderate storms at 46%.
- 3 Up to 84% of full halo CME driving storms originated close to the disk center; but only partial halo CMEs from outside  $\pm 45^\circ$  of the CMD were geoeffective.
- 4 Storms associated with FH CMEs were mostly intense storms (Mean  $Dst = -128$  nT); those associated with PH CMEs were moderate with avg  $Dst = -92$  nT.



# Main results

## Summary of main results

- 1 Geoeffective parameters ( $B_s$ ,  $V_{sw}$ ,  $V_{cmes}$ ) were of higher values for FH CMEs compared to those following PH CMEs.
- 2 FH CMEs were associated with class M flares on *avg*, class C on *avg* for PH CMEs.
- 3 26% of identified GMS were possibly driven by multiple (interacting) CMEs of which up to 69% were intense storms.
- 4 About half of all non-halo CME-driven storms were found in the declining phase of SC23

# Conclusion

# Conclusion

- 1 We have presented the results of a descriptive and statistical analysis of magnetic storms and associated solar and IP precursors during first 11-year of SC 23.

# Conclusion

- 1 We have presented the results of a descriptive and statistical analysis of magnetic storms and associated solar and IP precursors during first 11-year of SC 23.
- 2 The present results are significant for the long term space weather modelling and prediction.

# Conclusion

- 1 We have presented the results of a descriptive and statistical analysis of magnetic storms and associated solar and IP precursors during first 11-year of SC 23.
- 2 The present results are significant for the long term space weather modelling and prediction.
- 3 **Ref:** J. Uwamahoro and L. A. Mackinnell, *Solar and interplanetary precursors of geomagnetic storms in solar cycle 23*, ASR, Vol 51, pp.395– 410, 2013.

# Conclusion

- 1 We have presented the results of a descriptive and statistical analysis of magnetic storms and associated solar and IP precursors during first 11-year of SC 23.
- 2 The present results are significant for the long term space weather modelling and prediction.
- 3 **Ref:** J. Uwamahoro and L. A. Mackinnell, *Solar and interplanetary precursors of geomagnetic storms in solar cycle 23*, ASR, Vol 51, pp.395– 410, 2013.

1 **Thanks for your attention!**